

Wetlands Solutions *for the Future*

Learning from ALFAwetlands and
REWET Projects

 17 September 2026

 Brussels, Belgium



Funded by
the European Union

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or CINEA. Neither the European Union nor the granting authority can be held responsible for them.



ILMATIETEEN LAITOS
METEOROLOGISKA INSTITUTET
FINNISH METEOROLOGICAL INSTITUTE



RESEARCH INSTITUTE
NATURE AND FOREST

HARNESSING MICROBIAL COMMUNITIES FOR SUCCESSFUL WETLAND RESTORATION

Laura Kuusemets, Kaido Soosaar, Maarja Öpik, Mika Aurela, Arta Bārdule, Aldis Butlers, Romain Darnajoux, Tom De Dobbelaer, Laura Escarmena, Laure Gandois, Vincent Jassey, Jyrki Jauhiainen, Sari Juutinen, Rana Kanaan, Ain Kull, Tuula Larmola, Andis Lazdiņš, Ülo Mander, Maud Raman, José Miguel Sánchez Pérez, Sílvia Poblador, Francesc Sabater, Sabine Sauvage, Thomas Schindler, Bo Maxwell Stevens, Julien Tournebize, Liisa Ukonmaanaho, **Mikk Espenberg** (University of Tartu)



Funded by
the European Union

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or CINEA. Neither the European Union nor the granting authority can be held responsible for them.

Europe has lost 50–80% of historical **wetlands**

(Davidson, 2014;
Verhoeven, 2014)

Why wetlands matter?



Consequences:

- reduced **carbon** and **nitrogen** **storage**
- more **greenhouse** **gases**
- poorer **water** **quality** and **regulation**
- reduced **biodiversity**

(Mander, Öpik, Espenberg,
2025)

What happens when wetlands are drained?



Dry soil → oxygen increases → decomposition
→ GHG emissions

Wet soil → low oxygen → carbon/nitrogen stored

Why microbes matter?

Microbes (bacteria, archaea, fungi) control how **carbon and nutrients** move through **wetlands**.

They determine whether **wetlands store carbon** or **release greenhouse gases**.

Different microbial groups produce or consume these greenhouse gases

New Phytologist




Tansley insight | [Open Access](#)

Global peatland greenhouse gas dynamics: state of the art, processes, and perspectives

Ülo Mander ✉ Maarja Öpik, Mikk Espenberg

First published: 21 February 2025 | <https://doi.org/10.1111/nph.20430> | Cite this article as



Key topics and research perspectives	Importance	References
Methanogenesis vs methanotrophy	Improved understanding of the processes regulating CH ₄ balances, underpin strategies to reduce CH ₄ emissions as well as fluvial C losses	Gios <i>et al.</i> (2024); Espenberg <i>et al.</i> (2024)
Denitrification vs nitrification vs DNRA	Improved understanding of the processes regulating N ₂ O balances, underpin strategies to reduce gaseous emissions as well as fluvial N losses	Bösch <i>et al.</i> (2023); Espenberg <i>et al.</i> (2024); Masta <i>et al.</i> (2024)
Impact of BVOCs on GHG fluxes in peatlands	Largely unknown, initial studies indicate impact on CH ₄ fluxes	Junninen <i>et al.</i> (2022)
Integrated impact of soil moisture changes and microbiome dynamics on GHG fluxes in peatlands	Microbiome dynamics is a potential indicator for modelling of peatlands C and N	Bahram <i>et al.</i> (2022)
Impact of peat microbiome diversity and mycorrhiza on GHG fluxes in peatlands	Largely unknown, the impact on GHG fluxes yet to be fully analyzed	Storer <i>et al.</i> (2018); Wang <i>et al.</i> (2021); Page <i>et al.</i> (2022)

Microbial pathways controlling greenhouse gas fluxes are highly complex



Unraveling microbial processes involved in carbon and nitrogen cycling and greenhouse gas emissions in rewetted peatlands by molecular biology

Open access | Published: 16 March 2024

Volume 167, pages 609–629, (2024) | Cite this article


✓ You have full access to this open access article

Download PDF

Save article

Emilie Gios, Erik Verbruggen, Joachim Audet, Rachel Burns, Klaus Butterbach-Bahl, Mikk Espenberg, Christian Fritz, Stephan Glatzel, Gerald Jurasinski, Tuula Larmola, Ülo Mander, Claudia Nielsen, Andres F. Rodriguez, Clemens Scheer, Dominik Zak & Hanna M. Silvennoinen

6883 Accesses | 16 Citations | 5 Altmetric | Explore all metrics →



Gios et al. 2024, *Biogeochemistry*

Article | Open access | Published: 17 March 2022

Structure and function of the soil microbiome underlying N₂O emissions from global wetlands




Mohammad Bahram, Mikk Espenberg, Jaan Pärn, Laura Lehtovirta-Morley, Sten Anslan, Kuno Kasak,

Urmas Kõljalg, Jaan Liira, Martin Maddison, Mari Moora, Ülo Niinemets, Maarja Öpik, Meelis Pärtel, Kaido

Soosaar, Martin Zobel, Falk Hildebrand, Leho Tedersoo & Ülo Mander

Nature Communications 13, Article number: 1430 (2022) | Cite this article

37k Accesses | 227 Citations | 80 Altmetric | These authors contributed equally: Mohammad Bahram, Mikk Espenberg, Jaan Pärn, Leho Tedersoo, Ülo Mander.



Bahram et al. 2022, *Nature Communications*

One soil – many players – What do not we know yet?

1) Interactions. Are cycles and microbial domains interconnected?

Carbon, nitrogen, and metals cycles – **bacteria, archaea, and fungi.**

2) Restoration effects. Can restoration restore microbial processes?

Rewetting changes soil conditions – and microbial communities respond.

3) European-scale patterns. Do responses differ across wetland types?


Most studies are site-specific, limiting **generalisation across climates and land-use types.**

Study sites

- **33 sites**
- **6 countries** (Spain, France, Belgium, Latvia, Estonia, Finland)
- **8 wetland ecosystem types**
 - Unmanaged wetlands
 - Rewetted wetlands
 - Former peat extraction areas
 - Grazed peatlands
 - Reforested peatlands
 - Continuous cover forestry
 - Drained peatland forests
 - Treatment wetlands
- **543 soil samples**

Locations of the study sites.

Background map: European Wetland Map (Tegetmeyer *et al.*, 2024).



Methods: One of the largest wetland microbial studies across Europe


- **Soil sampling from three depths:**
 - 0-10 cm
 - 10-20 cm
 - 20-40 cm
- **Soil physicochemical parameters:**
 - Soil pH, TC, TN, NH_4 , NO_3
 - Soil water content (SWC)
 - K, Ca, Mg, Zn, Fe, Cr, Cd, Mn, Ni, P, Cu, Pb, Co, B, Al, Na, S
- **Microbial communities:** DNA extraction; sequencing of bacterial, archaeal, and fungal communities
- **GHG measurements:** CO_2 , CH_4 , N_2O



Soil sampling.

Microbial community diversity in Europe


- Peat extraction reduces diversity.
- Rewetting partially restores diversity.
- Diversity alone is not enough.



Panels show: (A) bacterial, (B) archaeal, (C) fungal Shannon diversity.

Different management practices support different prokaryotic (bacteria, archaea) functions


- **Natural and rewetted wetlands** support methane cycling → indicating **recovery of ecosystem functions**
- **Grazed peatlands** stimulate nutrient cycling → **driven by dung inputs.**
- **Treatment wetlands and reforested sites** → aligns with **high microbial diversity.**



Heatmap of predicted functional profiles for bacterial (A) and archaeal (B) communities based on functional annotation of prokaryotic taxa (FAPROTAX).

Plant communities strongly influence fungal communities and ecosystem functioning


- **Highest decomposition in open and grazed wetlands.**
- **Forested sites are mycorrhizal-dominated** → stronger plant–fungal coupling.
- **Grazed peatlands show dung saprotroph dominance** & high methanogenesis → dung inputs can boost CH₄ production.




Relative abundance (%) of fungal trophic guilds assigned using the FungalTraits database across the studied ecosystem types.

Microbial networks reveal restoration success

- Rewetted wetlands resemble natural wetlands
 - Network structure recovers after rewetting
 - Key microbial groups are linked with GHG fluxes
- Microbial groups linked with methane, nitrous oxide and carbon dioxide fluxes.




Networks of bacterial genera, soil properties, and GHG emissions across site types.




Networks of bacterial genera, soil properties, and GHG emissions across soil depths.

Metal availability strongly influences microbial communities and greenhouse gas processes

- **Iron (Fe), magnesium (Mg), and potassium (K)** are the strongest metal predictors of microbial taxa → have broad, cross-domain influences on wetland soil microbial communities.
- **Copper (Cu), nickel (Ni), and manganese (Mn)** show stronger domain-specific associations with bacterial, archaeal, and fungal communities, respectively.



Networks of fungal genera, soil properties, and GHG emissions across site types.



Networks of fungal genera, soil properties, and GHG emissions across soil depths.

Take-home message

- Rewetting restores microbial functioning
- Microbial networks indicate restoration success
- Metal availability influences greenhouse gas processes

What does this mean for wetland restoration?

- Rewetting restores key ecosystem processes
- Monitoring microbial networks can improve restoration assessment

THANK YOU

Contact:

Mikk Espenberg
mikk.espenberg@ut.ee



UT Environmental Microbiology Lab

